

Table 2

## Characteristics of major non-exploited brown coal deposits

	Name of deposit	Thickness of coal seams (in m)	Depth to the base	Overburden-to deposit thickness ratio	Calorific value (kcal/kg)	Ash content (%)	Mean. content of sulfur (%)
1	Babina - Żarki	10.7	140.0	-	2,229	18.28	1.10
2	Cybinka	16.6	94.0	5.6	2,236	18.40	1.41
3	Gubin	10.9	83.7	7.4	2,240	12.86	1.42
4	Głowaczów	4.8	37.1	6.5	1,820	28.56	0.42
5	Legnica p. Północ	22.0	207.2	8.7	2,194	18.72	1.42
6	Legnica p. Wschód	19.8	136.3	7.3	2,206	19.05	1.33
7	Legnica p. Zachód	21.0	158.8	6.6	2,371	20.10	0.76
8	Mosty	9.3	105.0	8.6	2,219	17.19	1.63
9	Piaski	6.1	48.5	7.3	2,082	24.80	1.44
10	Rogóżno	18.7	104.5	4.9	2,244	18.90	3.99
11	Rzepin	12.2	97.3	7.9	2,164	15.14	1.20
12	Sądów	12.2	127.5	10.2	2,196	18.80	1.38
13	Ścinawa	20.2	207.1	8.9	2,276	12.88	0.48
14	Torzym	21.4	180.8	7.9	2,270	16.80	1.81
15	Trzcianka	6.0	80.0	8.6	1,888	28.20	1.84
16	Złoczew	46.2	259.1	4.5	2,021	21.67	1.18