

Table 2

**CHARACTERISTICS OF MAJOR NON-EXPLOITED BROWN COAL DEPOSITS IN POLAND
as of 31.XII.2019**

	Name of deposit	Thickness of coal seams (m)	Depth to the base (m)	Overburden -to deposit thickness ratio	Calorific value (kcal/kg)	Ash content (%)	Mean content of sulfur (%)
1	Babina – Żarki	10.7	140.0	-	2,229	18.28	1.10
2	Bełchatów-pole Kamieńsk	18.1	184.6	9.2	1,944	25.81	1.02
3	Cybinka	8.8	78.0	7.9	2,251	17.16	1.32
4	Czempin	33.4	290.3	7.6	2,263	16.55	1.10
5	Dęby Szlacheckie	8.5	-	16.9	1,928	24.96	1.48
6	Gostyń	33.7	245.4	6.3	2,128	19.92	1.09
7	Gubin	7.6	99.7	9.3-16.0	2,240	20.27	2.69
8	Gubin 1	10.4	85.3	9.0	2,183	21.13	1.89
9	Gubin 2	13.3	127.6	8.2-8.7	2,254	17.86	3.71
10	Gubin-Zasieki-Brody	10.4	96.3	10.0	2,243	17.55	1.17
11	Głowaczów	4.8	37.1	6.5	1,820	28.56	0.42
12	Krzywin	23.6	221.9	11.9	2,229	14.89	0.70
13	Legnica-pole Północne	22.0	207.2	8.7	2,194	18.72	1.42
14	Legnica-pole Wschodnie	18.1	155.4	7.6	2,070	21.80	0.42
15	Legnica-pole Zachodnie	21.0	158.8	6.6	2,254	18.14	0.74
16	Lubsko	12.8	117.7	8.7	2,273	17.25	2.43
17	Łęki Szlacheckie	18.8	130.3	8.0	1,908	15.30	1.52
18	Mosina	32.1	225.1	6.6	2,210	18.70	0.66
19	Mosty	9.1	82.7	8.0	2,218	17.19	1.63
20	Nakło	18.5	-	7.3	1,983	24.13	0.90
21	Naramowice	28.0	212.5	6.9	2,076	19.40	0.50
22	Oczkowice	12.8	133.4	9.7	2,429	13.10	0.81
23	Piaski	6.1	48.5	7.3	2,072	24.90	1.44
24	Radomierzyce	17.6	78.5	4.5	1,868	17.58	1.12
25	Rogóżno	18.7	104.5	4.9	2,241	24.54	3.79
26	Ruja	16.8	140.4	8.0	2,237	18.57	0.71
27	Rzepin	12.2	97.3	7.9	2,164	15.14	1.20
28	Sądów	12.2	127.5	10.2	2,124	19.16	1.39
29	Szamotuły	23.1	179.6	7.6	2,340	13.40	0.40
30	Ścinawa	20.2	207.1	8.9	2,275	12.88	0.64
31	Torzym	21.4	180.8	7.9	2,270	16.80	1.81
32	Trzcianka	4.2	45.9	11.9	2,004	23.07	1.84
33	Więcbork	20.0	201.2	9.7	1,853	28.00	0.96
34	Złoczew	51.4	266.6	5.0	1,968	21.29	2.14